Subject:

CDC - Adults and Older Adult Adverse Drug Events - Medication Safety Program

<u>Skip directly to search Skip directly to A to Z list Skip directly to navigation Skip directly to site content Skip directly to page options</u>

CDC Home

Note: Javascript is disabled or is not supported by your browser. For this reason, some items on this page will be unavailable. For more information about this message, please visit this page: About CDC.gov.

Medication Safety Program

Medication Safety Program

Basics

Program Focus and Activities

For Parents: Young Children and Adverse Drug Events

Adults & Older Adult Adverse Drug Events

Campaigns and Initiatives

Resource Library

Publications by Year

Publications by Topic

The PROTECT Initiative

Partners/Members

Adults and Older Adult Adverse Drug Events

On this Page

Adverse drug events are a large public health problem

<u>Some medicines require blood testing and these are more likely to cause serious adverse events</u>
<u>Unintentional overdoses of pain relievers cause many deaths</u>

Some medicines may not be effective in improving your health

For More Information



Medicines cure infectious diseases, prevent problems from chronic diseases, and alleviate pain and suffering for millions of Americans every day. But medicines can also cause harm. When someone has

1

been harmed by a medicine, they have had an adverse drug event. To reduce the risk of harm from adverse drug events adults should:

Keep a list of your medicines
Follow directions
Ask questions
Keep up with any blood testing recommended by your doctor
Take pain relievers and antibiotics only as directed

Adverse drug events are a large public health problem.

Adverse drug events cause over 700,000 emergency department visits each year. Nearly 120,000 patients each year need to be hospitalized for further treatment after emergency visits for adverse drug events. As more and more people take more medicines, the risk of adverse events may increase.

As people age, they typically take more medicines. Older adults (65 years or older) are twice as likely as others to come to emergency departments for adverse drug events (over 177,000 emergency visits each year) and nearly seven times more likely to be hospitalized after an emergency visit.

Top of Page

Some medicines require blood testing and these are more likely to cause serious adverse events.

Some medicines need blood testing to help make sure the dose is just right for you. Over 40% of emergency visits which require patients to be hospitalized are caused by just a few of these medicines which require regular monitoring with blood tests.

Tip:

Ask if you need blood testing. Ask your doctor or pharmacist if you are taking any medicines that need blood testing and pay particular attention to taking these drugs properly and getting regular blood testing and follow-up. Common drugs that can require monitoring are

Blood thinners (e.g., <u>warfarin</u>)
Diabetes medicines (e.g., <u>insulin</u>)
Seizure medicines (e.g., <u>phenytoin</u>, <u>carbamazepine</u>)
Heart medicine (e.g., <u>digoxin</u>)

Top of Page

Unintentional overdoses of pain relievers cause many deaths

Medicines to relieve pain improve the quality of life for millions of Americans. However, in 2004, over 7,500 Americans died of unintentional overdoses of opioid analgesics (pain medicines such as methadone, oxycodone, and hydrocodone), more people than from cocaine or heroin

Tip:

Take pain relievers only as directed. If you are taking opioid pain relievers, be sure to tell your doctor about all other medicines you are taking because some medicines, when taken with pain relievers, can cause an overdose.

Top of Page

Some medicines may not be effective in improving your health

Antibiotics can kill bacteria but not viruses. Most colds, coughs, flu, sore throats, and runny noses are caused by viruses. Taking antibiotics for viral infections will not cure viral infections, keep others from catching the illness, or help you feel better. Although antibiotics are good drugs for certain types of infections, they are also the type of medicines that cause the most emergency visits for adverse drug events. Nearly 100,000 adults are treated in emergency departments each year because of adverse events from antibiotics.

Tip:

Do not ask for antibiotics when a doctor says they are not needed. They will not help treat viral infections.

If you are prescribed an antibiotic, make sure to take all the medicine as prescribed, even if you feel better. Do not save antibiotics "for later."

Top of Page

For More Information

For Safe Use of Oral Anticoagulants, SOS Rx

Unintentional Drug Poisoning in the United States, CDC

Emergency Department Visits Involving Nonmedical Use of Selected Prescription Drugs—United States, 2004–2008, CDC

Tips to Prevent Poisonings, CDC

Get Smart: Know When Antibiotics Work, CDC

Top of Page

Email page link

Print page

Get email updates

To receive email updates about this page, enter your email address:

Enter Email Address Submit Button [Submit] What's this?

Contact Us:

Centers for Disease Control and Prevention 1600 Clifton Rd Atlanta, GA 30333 800-CDC-INFO (800-232-4636) TTY: (888) 232-6348

Contact CDC-INFO



File Formats Help:



















How do I view different file formats (PDF, DOC, PPT, MPEG) on this site?

Email

Print

Page last reviewed: September 28, 2010 Page last updated: October 2, 2012

Content source: Centers for Disease Control and Prevention

National Center for Emerging and Zoonotic Infectious Diseases (NCEZID)

Division of Healthcare Quality Promotion (DHQP)

Home

A-Z Index

Site Map

Policies

Using this Site

Link to Us

All Languages

CDC Mobile

Contact CDC



Centers for Disease Control and Prevention $\,$ 1600 Clifton Rd. Atlanta, GA $\,$

30333, USA

800-CDC-INFO (800-232-4636) TTY: (888) 232-6348 - Contact CDC-INFO

A-Z Index

- 1. <u>A</u>
- 2. <u>B</u>
- 3. <u>C</u>
- 4. <u>D</u>
- 5. <u>E</u>
- 6. <u>F</u>
- 7. <u>G</u>
- 8. <u>H</u>
- 9. <u>I</u>
- 10. <u>J</u>
- 11. <u>K</u>
- 12. <u>L</u>
- 13. <u>M</u>
- 14. <u>N</u>
- 15. <u>O</u>
- 16. <u>P</u>
- 17. <u>Q</u>
- 18. <u>R</u>
- 19. <u>S</u>
- 20.<u>T</u>
- 21. <u>U</u>
- 22.<u>V</u>
- $23.\underline{W}$
- 24.<u>X</u>
- 25. <u>Y</u>
- 26.<u>Z</u>
- 27. <u>#</u>